

Dynamical Systems, Fractal Geometry and Diophantine Approximations

Carlos Gustavo Tamm de Araujo Moreira
 IMPA

December 13, 2017

Abstract

We describe in this survey several results relating Fractal Geometry, Dynamical Systems and Diophantine Approximations, including a description of recent results related to geometrical properties of the classical Markov and Lagrange spectra and generalizations in Dynamical Systems and Differential Geometry.

1 Introduction

The theory of Dynamical Systems is concerned with the asymptotic behaviour of systems which evolve over time, and gives models for many phenomena arising from natural sciences, as Meteorology and Celestial Mechanics. The study of a number of these models had a fundamental impact in the development of the mathematical theory of Dynamical Systems. An important initial stage of the theory of Dynamical Systems was the study by Poincare of the restricted three-body problem in Celestial Mechanics in late nineteenth century, during which he started to consider the qualitative theory of differential equations and proved results which are also basic to Ergodic Theory, as the famous Poincare's recurrence lemma. He also discovered during this work the *homoclinic* behaviour of certain orbits, which became very important in the study of the dynamics of a system. The existence of transverse homoclinic points implies that the dynamics is quite complicated, as remarked already by Poincare: "Rien n'est plus propre a nous donner une idee de la complication du probleme des trois corps et en general de tous les problemes de Dynamique...", in his classic *Les Methodes Nouvelles de la Mecanique Celeste* ([57]), written in late 19th century. This fact became clearer much decades later, as we will discuss below.

Poincare's original work on the subject was awarded a famous prize in honour of the 60th birthday of King Oscar II of Sweden. There is an interesting history related to this prize. In fact, there were two versions of Poincare's work presented for the prize, whose corresponding work was supposed to be published in the famous journal *Acta Mathematica* - a mistake was detected by the Swedish mathematician Phragmen in the first version. When Poincare became aware of that, he rewrote the paper, including the quotation above calling the attention to the great complexity of dynamical problems related to homoclinic

intersections. We refer to the excellent paper [61] by Jean-Christophe Yoccoz describing such an event.

We will focus our discussion of Dynamical Systems on the study of flows (autonomous ordinary differential equations) and iterations of diffeomorphisms, with emphasis in the second subject (many results for flows are similar to corresponding results for diffeomorphisms).

The first part of this work is related to the interface between Fractal Geometry and Dynamical Systems. We will give particular attention to results related to dynamical bifurcations, specially *homoclinic bifurcations*, perhaps the most important mechanism that creates complicated dynamical systems from simple ones. We will see how the study of the fractal geometry of hyperbolic sets has a central rôle in the study of dynamical bifurcations. We shall discuss recent results and ongoing works on fractal geometry of hyperbolic sets in arbitrary dimensions.

The second part is devoted to the study of the interface between Fractal Geometry and Diophantine Approximations. The main topic of this section will be the study of geometric properties of the classical Markov and Lagrange spectra - we will see how this study is related to the study of sums of regular Cantor sets, a topic which also appears naturally in the study of homoclinic bifurcations (which seems, at a first glance, to be a very distant subject from Diophantine Approximations).


The third part is related to the study of natural generalizations of the classical Markov and Lagrange spectra in Dynamical Systems and in Differential Geometry - for instance, we will discuss properties of generalized Markov and Lagrange spectra associated to geodesic flows in manifolds of negative curvature and to other hyperbolic dynamical systems. This is a subject with much recent activity, and several ongoing relevant works.

Acknowledgements: We would like to warmly thank Jacob Palis for very valuable discussions on this work.

2 Fractal Geometry and Dynamical Systems

2.1 Hyperbolic sets and Homoclinic Bifurcations

The notion of *hyperbolic systems* was introduced by Smale in the sixties, after a global example provided by Anosov, namely the diffeomorphism $f(x; y) = (2x + y; x + y) \pmod{1}$ of the torus $\mathbb{T}^2 = \mathbb{R}^2 / \mathbb{Z}^2$ and a famous example, given by Smale himself, of a *horseshoe*, that is a robust example of a dynamical system on the plane with a transverse homoclinic point as above, which implies a rich dynamics - in particular the existence of infinitely many periodic orbits. The figure below depicts a horseshoe.



(the dynamics sends the square $(ABCD)$ onto the domain bounded by $(A'B'C'D')$).

Let $\Lambda \subset M$ be a compact subset of a manifold M . We say that Λ is a hyperbolic set for a diffeomorphism $f : M \rightarrow M$ if $f(\Lambda) = \Lambda$ and there is a decomposition $T_\Lambda M = E^s \oplus E^u$ of the tangent bundle of M over Λ such that $Df^n|_{E^s}$ is uniformly contracting and $Df^n|_{E^u}$ is uniformly expanding. We say that f is hyperbolic if the limit set of its dynamics is a hyperbolic set.

It is important to notice that when a diffeomorphism has a transverse homoclinic point then its dynamics contains a non-trivial invariant hyperbolic set which is (equivalent to) a horseshoe - this explains the complicated situation discovered by Poincaré as mentioned above.

The importance of the notion of hyperbolicity is also related to the *stability conjecture* by Palis and Smale, according to which structurally stable dynamical systems are essentially the hyperbolic ones (i.e. the systems whose limit set is hyperbolic). After important contributions by Anosov, Smale, Palis, de Melo, Robbin and Robinson, this conjecture was proved in the C^1 topology by Mañé ([21]) for diffeomorphisms and, later, by Hayashi ([13]) for flows. The stability conjecture (namely the statement that structural stability implies hyperbolicity) is still open in the C^k topology for $k \geq 2$.


As mentioned in the introduction, homoclinic bifurcations are perhaps the most important mechanism that creates complicated dynamical systems from simple ones. This phenomenon takes place when an element of a family of dynamics (diffeomorphisms or flows) presents a hyperbolic periodic point whose stable and unstable manifolds have a non-transverse intersection. When we connect, through a family, a dynamics with no *homoclinic points* (namely, intersections of stable and unstable manifolds of a hyperbolic periodic point) to another one with a *transverse* homoclinic point (a homoclinic point where the intersection between the stable and unstable manifolds is transverse) by a family of dynamics, we often go through a homoclinic bifurcation.

Homoclinic bifurcations become important when going beyond the hyperbolic theory. In the late sixties, Sheldon Newhouse combined homoclinic bifurcations with the complexity already available in the hyperbolic theory and some new concepts in Fractal Geometry to obtain dynamical systems far more complicated than the hyperbolic ones. Ultimately this led to his famous result on the coexistence of infinitely many periodic attractors (which we will discuss later). Later on, Mora and Viana ([38]) proved that any surface diffeomorphism

morphism presenting a homoclinic tangency can be approximated by a diffeomorphism exhibiting a Henon-like strange attractor (and that such diffeomorphisms appear in any typical family going through a homoclinic bifurcation).

Palis conjectured that any diffeomorphism of a surface can be approximated arbitrarily well in the C^k topology by a hyperbolic diffeomorphism or by a diffeomorphism displaying a homoclinic tangency. This was proved by Pujals and Sambarino ([58]) in the C^1 topology. Palis also proposed a general version of this conjecture: any diffeomorphism (in arbitrary ambient dimension) can be approximated arbitrarily well in the C^k topology by a hyperbolic diffeomorphism, by a diffeomorphism displaying a homoclinic tangency or by a diffeomorphism displaying a heteroclinic cycle (a cycle given by intersections of stable and unstable manifolds of periodic points of different indexes). A major advance was done by Crovisier and Pujals ([6]), who proved that any diffeomorphism can be approximated arbitrarily well in the C^1 topology by a diffeomorphism displaying a homoclinic tangency, by a diffeomorphism displaying a heteroclinic cycle or by an *essentially hyperbolic* diffeomorphism: a diffeomorphism which displays a finite number of attractors whose union of basins is open and dense.

The first natural problem related to homoclinic bifurcations is the study of *homoclinic explosions* on surfaces: We consider one-parameter families (f_μ) , $\mu \in (-1; 1)$ of diffeomorphisms of a surface for which f_μ is uniformly hyperbolic for $\mu < 0$, and f_0 presents a quadratic homoclinic tangency associated to a hyperbolic periodic point (which may belong to a *horseshoe* - a compact, locally maximal, hyperbolic invariant set of saddle type). It unfolds for $\mu > 0$ creating locally two transverse intersections between the stable and unstable manifolds of (the continuation of) the periodic point. A main question is: what happens for (most) positive values of μ ? The following figure depicts such a situation for $\mu = 0$.



Fractal sets appear naturally in Dynamical Systems and fractal dimensions when we try to measure fractals. They are essential to describe most of the main results in this presentation.

Given a metric space X , it is often true that the minimum number $N(r)$ of balls of radius r needed to cover X is roughly proportional to $1/r^d$, for some positive constant d , when r becomes small. In this case, d will be the *box dimension* of X . More precisely, we

define the (upper) box dimension of X as

$$d(X) = \limsup_{r \rightarrow 0} \frac{\log N(r)}{\log r}.$$

The notion of *Hausdorff dimension* of a set is more subtle, but more useful. The main difference with the notion of box dimension is that, while the box dimension is related to coverings of X by small balls of equal radius, the Hausdorff dimension deals with arbitrary coverings of X by balls of small (but not necessarily equal) radius.

Given a countable covering U of X by balls, $U = (B(x_i; r_i))_{i \in \mathbb{N}}$, we define its norm $\|U\|$ as $\|U\| = \max_i r_i$ (where r_i is the radius of the ball $B(x_i; r_i)$). Given $s \geq 0$, we define $H_s(U) = \sum_{i \in \mathbb{N}} r_i^s$.

The Hausdorff s -measure of X is

$$H_s(X) = \lim_{\epsilon \rightarrow 0} \inf_{\|U\| < \epsilon} H_s(U):$$

One can show that there is a unique real number, the *Hausdorff dimension* of X , which we denote by $HD(X)$, such that $s < HD(X) \Rightarrow H_s(X) = +\infty$ and $s > HD(X) \Rightarrow H_s(X) = 0$ (so $HD(X)$ can be defined shortly as

$$HD(X) = \inf \{s > 0; \inf_{X \subset \cup B(x_n; r_n)} \sum r_n^s = 0\}:$$

For "well-behaved" sets X - in particular for regular Cantor sets in ambient dimension 1 and horseshoes in ambient dimension 2, the box and Hausdorff dimensions of X coincide.

Regular Cantor sets on the line play a fundamental role in dynamical systems and notably also in some problems in number theory. They are defined by expanding maps and have some kind of self-similarity property: small parts of them are diffeomorphic to big parts with uniformly bounded distortion (we will give a precise definition in a while). In both settings, dynamics and number theory, a key question is whether the arithmetic difference (see definition below) of two such sets has non-empty interior.

A horseshoe in a surface is locally diffeomorphic to the Cartesian product of two regular Cantor sets: the so-called *stable* and *unstable* Cantor sets K^s and K^u of Σ , given by intersections of Σ with local stable and unstable manifolds of some points of the horseshoe. The Hausdorff dimension of Σ , which is equal to the sum of the Hausdorff dimensions of K^s and K^u , plays a fundamental role in several results on homoclinic bifurcations associated to Σ .

From the dynamics side, in the eighties, Palis and Takens ([46], [47]) proved the following theorem about homoclinic bifurcations associated to a hyperbolic set:

Theorem 2.1. *Let $(f^j)_{j \geq 0}$, $f^j \in C^1(M; M)$ be a family of diffeomorphisms of a surface presenting a homoclinic explosion at $t = 0$ associated to a periodic point belonging to a horseshoe Σ . Assume that $HD(\Sigma) < 1$. Then*

$$\lim_{\epsilon \rightarrow 0} \frac{m(H \setminus [0; \epsilon])}{m(H)} = 1;$$

where $H := \{f^j > 0\}$ is uniformly hyperbolic.

2.2 Regular Cantor sets - a conjecture by Palis

A central fact used in the proof of the above theorem by Palis and Takens is that if K_1 and K_2 are regular Cantor sets on the real line such that the sum of their Hausdorff dimensions is smaller than one, then $K_1 - K_2 = \{x - y \mid x \in K_1, y \in K_2\}$ is a set of zero Lebesgue measure (indeed of Hausdorff dimension smaller than 1). On the occasion, looking for some kind of characterization property for this phenomenon, Palis conjectured (see [44], [45]) that for generic pairs of regular Cantor sets (K_1, K_2) of the real line either $K_1 - K_2$ has zero measure or else it contains an interval (the last case should correspond in homoclinic bifurcations to open sets of tangencies). A slightly stronger statement is that, if K_1 and K_2 are generic regular Cantor sets and the sum of their Hausdorff dimensions is bigger than 1, then $K_1 - K_2$ contains intervals.

Another motivation for the conjecture was Newhouse's work in the seventies, when he introduced the concept of thickness of a regular Cantor set, another fractal invariant associated to Cantor sets on the real line. It was used in [41] to exhibit open sets of diffeomorphisms with persistent homoclinic tangencies, therefore with no hyperbolicity. It is possible ([42]) to prove that, under a dissipation hypothesis, in such an open set there is a residual set of diffeomorphisms which present in finitely many coexisting sinks. In [43], it is proved that under generic hypotheses every family of surface diffeomorphisms that unfold a homoclinic tangency goes through such an open set. It is to be noted that in the case described above with $HD(K) < 1$ (as studied in [47]) these sets have zero density. See [48] for a detailed presentation of these results. An important related question by Palis is whether the sets of parameter values corresponding to finitely many coexisting sinks have typically zero Lebesgue measure.

An earlier and totally independent development had taken place in number theory. In 1947, M. Hall ([12]) proved that any real number can be written as the sum of two numbers whose continued fractions coefficients (of positive index) are bounded by 4. More precisely, if $C(4)$ is the regular Cantor set (see general definition below) formed of such numbers in $[0; 1]$, then one has $C(4) + C(4) = [0; 2] \setminus \{1/4, 3/4, \dots\}$. We will discuss generalizations and consequences of this result in the next section.

A Cantor set $K \subset \mathbb{R}$ is a C^k -regular Cantor set, $k \geq 1$, if:

- i) there are disjoint compact intervals I_1, I_2, \dots, I_r such that $K \subset \bigcup_{j=1}^r I_j$ and the boundary of each I_j is contained in K ;
- ii) there is a C^k expanding map f defined in a neighbourhood of $\bigcup_{j=1}^r I_j$ such that, for each j , $f(I_j)$ is the convex hull of a finite union of some of these intervals I_s . Moreover, we suppose that f satisfies:
 - ii.1) for each j , $1 \leq j \leq r$ and n sufficiently big, $f^n(K \setminus I_j) = K$;
 - ii.2) $K = \bigcap_{n \in \mathbb{N}} f^{-n}(\bigcup_{j=1}^r I_j)$.

Remark 2.2. If k is not an integer, say $k = m + \alpha$, with $m \geq 1$ integer and $0 < \alpha < 1$ we assume that f is C^m and $f^{(m)}$ is α -Hölder.

We say that $f: I_1; I_2; \dots; I_r$ is a *Markov partition* for K and that K is *defined* by f .

Remark 2.3. In general, we say that a set $X \subset \mathbb{R}$ is a Cantor set if X is compact, without isolated points and with empty interior. Cantor sets in \mathbb{R} are homeomorphic to the classical ternary Cantor set $K_{1=3}$ of the elements of $[0; 1]$ which can be written in base 3 using only digits 0 and 2. The set $K_{1=3}$ is itself a regular Cantor set, defined by the map $f: [0; 1=3] \rightarrow [2=3; 1] \subset \mathbb{R}$ given by $f(x) = 3x - b3xc$.

An *interval of the construction of the regular Cantor set K* is a connected component of $f^{-n}(I_j)$ for some $n \in \mathbb{N}$, $j = 1, \dots, r$.

Given $s \in [1; k]$ and another regular Cantor set K , we say that K is close to K in the C^s topology if K has a Markov partition $f: t_1; t_2; \dots; t_r$ such that the interval t_j has endpoints close to the endpoints of I_j , for $1 \leq j \leq r$ and K is defined by a C^s map f which is close to f in the C^s topology.

The C^{1+} -topology is such that a sequence K_n converges to K if there is some $\epsilon > 0$ such that K_n is $C^{1+\epsilon}$ for every $n \geq 1$ and K_n converges to K in the C^{1+} -topology.

The concept of stable intersection of two regular Cantor sets was introduced in [25]: two Cantor sets K_1 and K_2 have stable intersection if there is a neighbourhood V of $(K_1; K_2)$ in the set of pairs of C^{1+} -regular Cantor sets such that $(\tilde{K}_1; \tilde{K}_2) \in V \Rightarrow \tilde{K}_1 \cap \tilde{K}_2 \neq \emptyset$.

In the same paper conditions based on renormalizations were introduced for Cantor



Jean-Christophe Yoccoz

The same result works if we replace stable intersection by d -stable intersection, which is defined by asking that any pair $(\tilde{K}_1; \tilde{K}_2)$ in some neighbourhood of $(K_1; K_2)$ satisfies $HD(\tilde{K}_1 \setminus \tilde{K}_2) \leq d$: most pairs of Cantor sets $(K_1; K_2) \in \mathcal{C}^\infty$ have d -stable intersection for any $d < HD(K_1) + HD(K_2) - 1$.

The open set U mentioned in the above theorem is very large in \mathcal{C}^∞ in the sense that generic n -parameter families in \mathcal{C}^∞ are actually contained in U .

The proof of this theorem depends on a sufficient condition for the existence of stable intersections of two Cantor sets, related to a notion of renormalization, based on the fact that small parts of regular Cantor sets are diffeomorphic to the whole set: the existence of a *recurrent compact set* of relative positions of limit geometries of them. Roughly speaking, it is a compact set of relative positions of regular Cantor sets such that, for any relative position in such a set, there is a pair of (small) intervals of the construction of the Cantor sets such that the renormalizations of the Cantor sets associated to these intervals belong to the interior of the same compact set of relative positions.

The main result is reduced to prove the existence of recurrent compact sets of relative positions for most pairs of regular Cantor sets whose sum of Hausdorff dimensions is larger than one. A central argument in the proof of this fact is a probabilistic argument à la Erdős: we construct a family of perturbations with a large number of parameters and show the existence of such a compact recurrent set with large probability in the parameter space (without exhibiting a specific perturbation which works).

Another important ingredient in the proof is the Scale Recurrence Lemma, which, under mild conditions on the Cantor sets (namely that at least one of them is not essentially affine), there is a recurrent compact set for renormalizations at the level of relative scales of limit geometries of the Cantor sets. This lemma is the fundamental tool in the paper [28], in which it is proved that, under the same hypothesis above, if K and K' are C^2 -regular Cantor sets, then $HD(K + K') = \min\{1; HD(K) + HD(K')\}$.

An important result in fractal geometry which is used in [39] is the famous Marstrand's theorem ([23]), according to which, given a Borel set $X \subset \mathbb{R}^2$ with $HD(X) > 1$ then, for almost every $\alpha \in \mathbb{R}$, $\alpha(X)$ has positive Lebesgue measure, where $\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}$ is given by $\alpha(x; y) = x - \alpha y$. In particular, if K_1 and K_2 are regular Cantor sets with $HD(K_1) + HD(K_2) > 1$ then, for almost every $\alpha \in \mathbb{R}$, $K_1 - \alpha K_2$ has positive Lebesgue measure. Moreira and Yuri Lima gave combinatorial alternative proofs of Marstrand's theorem, first in the case of Cartesian products regular Cantor sets ([19]) and then in the general case ([20]).

In [40], Moreira and Yoccoz proved the following fact concerning generic homoclinic bifurcations associated to two dimensional saddle-type hyperbolic sets (horseshoes) with Hausdorff dimension bigger than one: typically there are translations of the stable and unstable Cantor sets having stable intersection, and so it yields open sets of stable tangencies in the parameter line with positive density at the initial bifurcation value. Moreover, the union of such a set with the hyperbolicity set in the parameter line generically has full density at the initial bifurcation value. This extends the results of [52].

The situation is quite different in the C^1 -topology, in which stable intersections do not exist:

Theorem 2.5 ([26]). *Given any pair $(K; K')$ of regular Cantor sets, we can find, arbitrarily close to it in the C^1 topology, pairs $(\mathcal{K}; \mathcal{K}')$ of regular Cantor sets with $\mathcal{K} \setminus \mathcal{K}' = \emptyset$.*

Moreover, for generic pairs $(K; K')$ of C^1 -regular Cantor sets, the arithmetic difference $K - K'$ has empty interior (and so is a Cantor set).

The main technical difference between the C^1 case and the C^2 (or even C^{1+}) cases is the lack of bounded distortion of the iterates of α in the C^1 case, and this fact is fundamental for the proof of the previous result.

The previous result may be used to show that there are no C^1 robust tangencies between leaves of the stable and unstable foliations of respectively two given hyperbolic horseshoes $\mathcal{H}_1; \mathcal{H}_2$ of a diffeomorphism of a surface. This is also very different from the situation in the C^∞ topology - for instance, in [40] it is proved that, in the unfolding of a homoclinic or heteroclinic tangency associated to two horseshoes, when the sum of the correspondent stable and unstable Hausdorff dimensions is larger than one, there are generically stable tangencies associated to these two horseshoes. This result is done in the following

Theorem 2.6 ([26]). *Given a C^1 diffeomorphism f of a surface M having two (non necessarily disjoint) horseshoes $\mathcal{H}_1; \mathcal{H}_2$, we can find, arbitrarily close to it in the C^1 topology, a diffeomorphism \tilde{f} of the surface for which the horseshoes $\mathcal{H}_1; \mathcal{H}_2$ have hyperbolic continuations $\tilde{\mathcal{H}}_1; \tilde{\mathcal{H}}_2$, and there are no tangencies between leaves of the stable and unstable*

foliations of $\tilde{\nu}_1$ and $\tilde{\nu}_2$, respectively. Moreover, there is a generic set R of C^1 diffeomorphisms of M such that, for every $f \in R$, there are no tangencies between leaves of the stable and unstable foliations of $\nu_1; \nu_2$, for any horseshoes $\nu_1; \nu_2$ of f .

Since stable intersections of Cantor sets are the main known obstructions to density of hyperbolicity for diffeomorphisms of surfaces, the previous result gives some hope of proving density of hyperbolicity in the C^1 topology for diffeomorphisms of surfaces, a well-known question by Smale. In particular in the work [32] on a family of two-dimensional maps (the so-called Benedicks-Carleson *toy model*

so g_t has no attractors nor repellers in W . Most small values of t , and thus most g_t , here means that t is taken in a set of parameter values with Lebesgue density one at $t = 0$.

The construction of non-uniformly hyperbolic horseshoes for most parameters is a highly non-trivial counterpart of Yoccoz' proof [62] (based on the so-called *Yoccoz puzzles*) of the celebrated Jakobson's theorem in the context of heteroclinic explosions.

Of course, Palis and Yoccoz do not consider their result as the end of the line. Indeed, they expected the same results to be true for all cases $0 < HD(\) < 2$. However, to achieve that, it seems that their methods need to be considerably sharpened: it would be necessary to study deeper the dynamical recurrence of points near tangencies of higher order (cubic, quartic...) between stable and unstable curves. They also expected their results to be true in higher dimensions.

Finally, they hoped that the ideas introduced in that work might be useful in broader contexts. In the horizon lies a famous question concerning the standard family of area preserving maps (which is the family $(f_k)_{k \in \mathbb{R}}$ of diffeomorphisms of the torus $\mathbb{T}^2 = \mathbb{R}^2 / \mathbb{Z}^2$ given by $f_k(x; y) = (y + 2x + k \sin(2x); x) \pmod{1}$): can we find sets of positive Lebesgue measure in the parameter space such that the corresponding maps display non-zero Lyapunov exponents in sets of positive Lebesgue probability in phase space?

In space?

tions are inspired in the *asymptotic dimensions*, introduced by Falconer. We have analogous definitions for upper unstable dimensions. We prove the following results about these dimensions: given $1 \leq r \leq \infty$ and $\epsilon > 0$ there is a δ small C^∞ perturbation of the original diffeomorphism for which the hyperbolic continuation of f has a subhorseshoe \tilde{h} which has strong-stable foliations of codimensions j for $1 \leq j \leq r$ and which satisfies $\bar{d}_s^{(r)}(\tilde{h}) > \bar{d}_s^{(r)}(f) - \epsilon$.

In the second work in progress, in collaboration with W. Silva (which extends a previous joint work in codimension 1 - see [37]), we prove that if a horseshoe h has strong stable foliations of codimensions j for $1 \leq j \leq r$ and satisfies $\hat{d}_s(h) > r$ (which is equivalent to $\bar{d}_s^{(r)}(h) > r$) then it has a small C^∞ perturbation which contains a blender of codimension r : in particular C^1 images of stable Cantor sets of it (of the type $\tilde{h} \setminus W^s(x)$) in \mathbb{R}^r will typically have persistently non-empty interior. We also expect to prove that the Hausdorff dimension of these stable Cantor sets typically coincide with $\hat{d}_s(h)$, and this dimension depends continuously on h on these assumptions, which would imply typical continuity of Hausdorff dimensions of stable and unstable Cantor sets of horseshoes.

Let us recall two main ingredients in the proof by Moreira and Yoccoz of Palis' conjecture:

A recurrent compact set criterion for stable intersections (which implies that arithmetic differences persistently contain intervals).

An application of Erdős probabilistic method: a family of C^∞ small perturbations of a regular Cantor set (the second Cantor set is fixed) with a large number of parameters such that for most parameters there is a recurrent compact set for the corresponding pair of Cantor sets.

A variation of these ingredients is present in the proof of these results in collaboration of W. Silva: we develop a recurrent compact set criterion which implies that a given horseshoe (with a strong-stable foliation of codimension r) is a r -codimensional blender, and we prove that, if a horseshoe has a strong-stable foliation of codimension r and satisfies $\hat{d}_s(h) > r$ (which is equivalent to $\bar{d}_s^{(r)}(h) > r$), then there is a small C^∞ perturbation of it which has a recurrent compact set. In order to do this, we use the probabilistic method: we construct a family of perturbations with a large number of parameters, and show that for most parameters there is such a recurrent compact set.

An important geometrical tool in the proof of these results is the following generalization of Marstrand's theorem, which was proved in works by Lopez, Moreira, Romana and Silva:

Theorem 2.8. *Let X be a compact metric space, $(\Omega; P)$ a probability space and $\pi : X \rightarrow \mathbb{R}^r$ a measurable function. Informally, one can think of $\pi(\cdot) = (\cdot; \cdot)$ as a family of projections parameterized by \cdot . We assume that for some positive real numbers ϵ and C the following transversality property is satisfied:*

$$P[\exists \epsilon > 0 : d(\pi(x_1); \pi(x_2)) \geq \epsilon \implies d(x_1; x_2) \geq C \epsilon^r] = 1 \quad (1)$$

for all $\epsilon > 0$ and all $x_1, x_2 \in X$. Assume that $\dim X > r$. Then $\text{Leb}(\pi(X)) > 0$ for a.e. $\cdot \in \Omega$ and $\int_\Omega \text{Leb}(\pi(X))^{-1} dP < +\infty$.

3 Fractal Geometry and Diophantine Approximations

3.1 The classical Markov and Lagrange spectra

The results discussed in the previous section on regular Cantor sets have, somewhat surprisingly, deep consequences in number theory, which we discuss below. We begin by introducing the classical Lagrange spectrum.

Let α be an irrational number. According to Dirichlet's theorem, the inequality $|j - \frac{p}{q}| < \frac{1}{q^2}$ has in nitely many rational solutions $\frac{p}{q}$. Hurwitz improved this result by proving that $|j - \frac{p}{q}| < \frac{1}{\sqrt{5}q^2}$ also has in nitely many rational solutions $\frac{p}{q}$ for any irrational α , and that $\frac{1}{\sqrt{5}}$ is the largest constant that works for any irrational α . However, for particular values of α we can improve this constant.

More precisely, we define $k(\alpha) := \sup\{k > 0 \mid |j - \frac{p}{q}| < \frac{1}{kq^2} \text{ has in nitely many rational solutions}\}$. We have $k(\alpha) \geq \frac{1}{\sqrt{5}}$, $k(\alpha) \in \mathbb{R} \setminus \mathbb{Q}$ and $k(\frac{1+\sqrt{5}}{2}) = \frac{1}{\sqrt{5}}$. We will consider the set $L = \{k(\alpha) \mid \alpha \in \mathbb{R} \setminus \mathbb{Q}, k(\alpha) < +\infty\}$.

This set is called the Lagrange spectrum. Hurwitz's theorem determines the smallest element of L , which is $\frac{1}{\sqrt{5}}$. This set L encodes many diophantine approximations of real numbers. It is a classical subject the study of the geometric structure of L . Markov proved in 1879 ([24]) that

$$L \setminus (1; 3) = \{k_1 = \frac{1}{\sqrt{5}} < k_2 = \frac{1}{\sqrt{2}} < k_3 = \frac{1}{\sqrt{221}} < \dots\}$$

where k_n is a sequence (of irrational numbers whose squares are rational) converging to 3.

The elements of the Lagrange spectrum which are smaller than 3 are exactly the numbers of the form $\sqrt{9 - \frac{4}{z^2}}$ where z is a positive integer for which there are other positive integers x, y such that $1/x + 1/y + 1/z$ and (x, y, z) is a solution of the *Markov equation*

$$x^2 + y^2 + z^2 = 3xyz$$

Since this is a quadratic equation in x (resp. in y), whose sum of roots is $3yz$ (resp. $3xz$), given a solution (x, y, z) with $x < y < z$, we also have the two other solutions $(y, z; 3yz - x)$ and $(x, z; 3xz - y)$ - it is possible to prove that all solutions of the Markov equation appear in the following tree:



An important open problem related to Markov's equation is the *Unicity Problem*, formulated by Frobenius about 100 years ago: for any positive integers x_1, x_2, y_1, y_2, z with $x_1 + y_1 = z$ and $x_2 + y_2 = z$ such that (x_1, y_1, z) and (x_2, y_2, z) are solutions of Markov's equation we always have $(x_1, y_1) = (x_2, y_2)$?

If the Unicity Problem has an affirmative answer then, for every real $t < 3$, its pre-image $k^{-1}(t)$ by the function k above consists of a single $GL_2(\mathbb{Z})$ -equivalence class (this equivalence relation is such that

$$\frac{a + b}{c + d}; (a, b, c, d \in \mathbb{Z}; jad - bcj = 1.)$$


Despite the "beginning" of the set L is discrete, this is not true for the whole set L . As we mentioned in the introduction, M. Hall proved in 1947 ([12]) that if $C(4)$ is the regular Cantor set formed by the numbers in $[0; 1]$ whose coefficients in the continued fractions expansion are bounded by 4, then one has $C(4) + C(4) = [\sqrt{2} - 1; 4(\sqrt{2} - 1)]$. This implies that L contains a whole half line (for instance $[6; +\infty)$), and G. Freiman determined in 1975 ([9]) the biggest half line that is contained in L , which is $[c_F; +\infty)$, with

$$c_F = \frac{2221564096 + 283748\sqrt{462}}{491993569} = 4.52782956616 \dots$$

These last two results are based on the study of sums of regular Cantor sets, whose relationship with the Lagrange spectrum will be explained below.

Sets of real numbers whose continued fraction representation has bounded coefficients with some combinatorial constraints, as $C(4)$, are often regular Cantor sets, which we call Gauss-Cantor sets (since they are defined by restrictions of the Gauss map $g(x) = \frac{1}{x} - \text{int}(\frac{1}{x})$ from $(0; 1)$ to $[0; 1)$ to some convenient union of intervals).

We represent below the graphics of the Gauss map $g(x) = \frac{1}{x} - \text{int}(\frac{1}{x})$.



If the continued fraction of x is

$$x = [a_0; a_1; a_2; \dots] \stackrel{\text{def}}{=} a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$$

then we have the following formula for $k(\alpha)$:

$$k(\alpha) = \limsup_{n \rightarrow \infty} (p_n + q_n);$$

where $p_n = [a_n; a_{n+1}; a_{n+2}; \dots]$ and $q_n = [0; a_{n-1}; a_{n-2}; \dots; a_1]$.

The previous formula follows from the equality

$$j \frac{p_n}{q_n} = \frac{1}{(p_{n+1} + p_{n+1})q_n^2}; \quad \forall n \in \mathbb{N};$$

where

$$p_n = q_n = [a_0; a_1; a_2; \dots; a_n] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots + \frac{1}{a_n}}}; \quad n \in \mathbb{N}$$

are the convergents of the continued fraction of α .

There are many results which relate the dynamics of the Gauss map with the behaviour of continued fractions. For instance, the Khintchine-Levy theorem, which follows from techniques of Ergodic Theory, states that, for (Lebesgue) almost every $\alpha \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \frac{p_n}{q_n} = e^{-2} = 12 \log 2.$$

Remark: The following elementary general facts on Diophantine approximations of real numbers show that the best rational approximations of a given real number are given by convergents of its continued fraction representation:

For every $n \in \mathbb{N}$,

$$\left| \frac{p_n}{q_n} - \alpha \right| < \frac{1}{2q_n^2} \quad \text{or} \quad \left| \frac{p_{n+1}}{q_{n+1}} - \alpha \right| < \frac{1}{2q_{n+1}^2}$$

(moreover, for every positive integer n , there is $k \geq n-1; n; n+1$ with $j \frac{p_k}{q_k} < \frac{1}{\sqrt{5}q_n^2}$).

If $\left| \frac{p}{q} - \alpha \right| < \frac{1}{2q^2}$ then $\frac{p}{q}$ is a convergent of the continued fraction of α .

This formula for $k(\alpha)$ implies that we have the following alternative definition of the Lagrange spectrum L :

Let $\Sigma = (\mathbb{N}^*)^{\mathbb{Z}}$ be the set of all bi-infinite sequences of positive integers. If $\alpha = (a_n)_{n \in \mathbb{Z}} \in \Sigma$, let $p_n = [a_n; a_{n+1}; a_{n+2}; \dots]$ and $q_n = [0; a_{n-1}; a_{n-2}; \dots]; \forall n \in \mathbb{Z}$. We define $f(\alpha) = p_0 + p_0 = [a_0; a_1; a_2; \dots] + [0; a_{-1}; a_{-2}; \dots]$. We have

$$L = \infsup_{n \rightarrow \infty} f(\sigma^n \alpha); \alpha \in \Sigma;$$

where $\sigma: \Sigma \rightarrow \Sigma$ is the shift defined by $(\sigma(a_n)_{n \in \mathbb{Z}}) = (a_{n+1})_{n \in \mathbb{Z}}$.

Let us define the Markov spectrum M by

$$M = \infsup_{n \in \mathbb{Z}} f(\sigma^n \alpha); \alpha \in \Sigma;$$

It also has an arithmetical interpretation, namely

$$M = f\left(\inf_{(x,y) \in \mathbb{Z}^2 \setminus (0,0)} |f(x,y)|\right)^{-1}; \quad f(x,y) = ax^2 + bxy + cy^2; \quad b^2 - 4ac = 1g:$$

It is well-known (see [3]) that M and L are closed sets of the real line and $L \subset M$. We have the following result about the Markov and Lagrange spectra:

Theorem 3.1. ([27]; see also [22]) *Given $t \in \mathbb{R}$ we have*

$$HD(L \setminus (-t; t)) = HD(M \setminus (-t; t)) =: d(t)$$

and $d(t)$ is a continuous surjective function from \mathbb{R} to $[0;1]$. Moreover:

i) $d(t) = \min\{1; 2D(t)g\}$, where $D(t) := HD(k^{-1}(-t; t)) = HD(k^{-1}(t; t))$ is a continuous function from \mathbb{R} to $[0;1]$.

ii) $\max_{t \in \mathbb{R}} d(t) = 0g = 3$

iii) $d(\sqrt{12}) = 1$.

A fundamental tool in the proof of this result is the theorem below.

We say that a C^2 -regular Cantor set on the real line is *essentially a λ -ne* if there is a C^2 change of coordinates for which the dynamics that defines the corresponding Cantor set has zero second derivative on all points of that Cantor set. Typical C^2 -regular Cantor sets are not essentially a λ -ne.

The *scale recurrence lemma*, which is the main technical lemma of [39], can be used in order to prove the following

Theorem 3.2. ([28]) *If K and K' are regular Cantor sets of class C^2 and K is non essentially a λ -ne, then $HD(K + K') = \min\{HD(K) + HD(K'); 1g\}$:*

Remark 3.3. There is a presentation of a version of this result (with a slightly different hypothesis) in [60]. That version is also proved by Hochman and Shmerkin in [15].

The results of Markov, Hall and Freiman mentioned above imply that the Lagrange and Markov spectra coincide below 3 and above c_F . Nevertheless, Freiman ([7]) showed in 1968 that $M \cap L \neq \emptyset$; by exhibiting a number $\alpha \approx 3.1181 \in M \cap L$.

In 1973, Freiman [8] showed that

$$\alpha_\infty := \alpha_0(A_\infty) := [2; \overline{1_2; 2_3; 1; 2}] + [0; 1; 2_3; 1_2; 2; 1; \overline{2}] \in M \cap L$$

In a similar vein, Theorem 4 in Chapter 3 of Cusick-Flahive book [3] asserts that

$$\alpha_n := \alpha_0(A_n) := [2; \overline{1_2; 2_3; 1; 2}] + [0; 1; 2_3; 1_2; 2; 1; 2_n; \overline{1; 2; 1_2; 2_3}] \in M \cap L$$

for all $n \geq 4$. In particular, α_∞ is not isolated in $M \cap L$.

In collaboration with C. Matheus, we proved the following results about $M \cap L$:

Let X be the Cantor set

$$X := f[0; 1] \subset f[1; 2g^{\mathbb{N}} \text{ not containing the subwords in } Pg \quad (2)$$

where

$$P := f21212;2121_3;1_3212;12121_2;1_22121;2_3121_22_21;1_221_2212_3;1_23121_22_2;2_21_2212_31g$$

Also, let

$$b_\infty := [2; \overline{1_2; 2_3; 1; 2}] + [0; \overline{1; 2_3; 1_2; 2}] = \frac{P_{18229}}{41} = 3.2930442439 \dots$$

and

$$\begin{aligned} B_\infty &:= [2; 1; \overline{1; 2_3; 1; 2; 1_2; 2; 1_2; 2}] + [0; 1; 2_3; 1_2; 2; 1; 2_3; 1_2; 2; 1; 2_2; \overline{1; 2_3; 1; 2; 1_2; 2; 1_2; 2}] \\ &= 3.2930444814 \dots \end{aligned}$$

Theorem 3.4. ([30]) $fb_\infty; B_\infty g \in L$, $L \setminus (b_\infty; B_\infty) = \emptyset$; and $HD((M \cap L) \setminus (b_\infty; B_\infty)) = HD(X)$.

We implemented the algorithm of Jenkinson-Pollicott (see [17]) and we obtained the heuristic approximation $HD(X) = 0.4816 \dots$. We also exhibited a Cantor set $K(f1; 2; 2g) \subset X$ whose Hausdorff dimension can be easily (and rigorously) estimated as $0.353 < HD(K(f1; 2; 2g)) < 0.35792$ via some classical arguments explained in Palis-Takens book [47].

By exploiting the arguments establishing the above Theorem, we are able to exhibit new numbers in $M \cap L$, including a constant $c \in M \cap L$ with $c > 4$:

Proposition 3.5. The largest element of $(M \cap L) \setminus (b_\infty; B_\infty)$ is

$$c = \frac{77 + \frac{P_{18229}}{41}}{82} + \frac{17633692 \frac{P_{151905}}{41}}{24923467} = 3.29304447990138 \dots$$

To our best knowledge, c is the largest known element of $M \cap L$.

We also proved ([31]) that $M \cap L$ doesn't have full Hausdorff dimension:

Theorem 3.6. $HD(M \cap L) < 0.986927$.

One can get better heuristic bounds for $HD(M \cap L)$ thanks to the several methods in the literature to numerically approximate the Hausdorff dimension of Cantor sets of numbers with prescribed restrictions of their continued fraction expansions. By implementing the "thermodynamical method" introduced by Jenkinson-Pollicott in [16], we obtained the heuristic bound $HD(M \cap L) < 0.888$.

Our proof of Theorem 3.6 relies on the control of several portions of $M \cap L$ in terms of the sum-set of a Cantor set associated to continued fraction expansions prescribed by a "symmetric block" and a Cantor set of irrational numbers whose continued fraction expansions live in the "gaps" of a "symmetric block". As it turns out, such a control is possible thanks to our key technical Lemma saying that a sufficiently large Markov value given by the sum of two continued fraction expansions systematically meeting a "symmetric block" must belong to the Lagrange spectrum.

It follows that $M \cap L$ has empty interior, and so, since M and L are closed subsets of \mathbb{R} , $\overline{int(M)} = \overline{int(L)} \subset L \subset M$. In particular, we have the following

Corollary 3.7. $int(M) = int(L)$.

As a consequence, we recover the fact, proved in [9], that the biggest half-line contained in M coincides with the biggest half-line $[c_F; \infty)$ contained in L .

3.2 Other results on the fractal geometry of Diophantine approximations

In collaboration with Y. Bugeaud ([1]), we proved some results on sets of exact approximation order by rational numbers:

For a function $\psi : (0; +\infty) \rightarrow (0; +\infty)$, let

$$K(\psi) := \left\{ x \in \mathbb{R} : \left| \frac{p}{q} - x \right| < \psi(q) \text{ for infinitely many rational numbers } \frac{p}{q} \right\}$$

denote the set of ψ -approximable real numbers and let

$$\text{Exact}(\psi) := K(\psi) \cap \bigcup_{m \geq 2} K((1 - 1/m)\psi)$$

be the set of real numbers approximable to order ψ and to no better order.

The *lower order at infinity* $\omega(\psi)$ of a function $\psi : (0; +\infty) \rightarrow (0; +\infty)$ is defined by

$$\omega(\psi) = \liminf_{x \rightarrow +\infty} \frac{\log \psi(x)}{\log x}.$$

We say that a function $\psi : (0; +\infty) \rightarrow (0; +\infty)$ satisfies assumption (A) if $\psi(x) = o(x^{-2})$ and there exist real numbers c, ϵ and n_0 with $1 - \epsilon < 4$ such that, if the positive integers m, n satisfy $m > n \geq n_0$, then $\psi(m)m^c < \epsilon \psi(n)n^c$. We emphasize that the real number c occurring in (A) may be negative.

Theorem 3.8. *Let $\psi : (0; +\infty) \rightarrow (0; +\infty)$ be a function satisfying assumption (A) . Then the set $\text{Exact}(\psi)$ is uncountable.*

Theorem 3.9. *Let $\psi : (0; +\infty) \rightarrow (0; +\infty)$ be a function satisfying assumption (A) . If ω denotes the lower order at infinity of the function ψ , then*

$$\dim \text{Exact}(\psi) = \dim K(\psi) = \frac{2}{\omega}.$$

In another work collaboration with Y. Bugeaud ([2]), we proved that there are no typical real numbers from the point of view of Diophantine approximations, in a sense that we describe in what follows. Let ψ be an application from the set of positive integers into the set of nonnegative real numbers. Khintchine established that, if the function $q \mapsto q^2 \psi(q)$ is non-increasing and the series $\sum_{q \geq 1} q^{-2} \psi(q)$ diverges, then the set $K(\psi)$ has full Lebesgue measure (Beresnevich, Dickinson and Velani proved later the same result assuming that ψ is just non-increasing). We show that, for almost every real number x , there is a function ψ which satisfies good "regularity" conditions (on the speed of decreasing of ψ) - for instance $q \mapsto q^2 \psi(q)$ is non-increasing, such that the series $\sum_{q \geq 1} q^{-2} \psi(q)$ diverges but the inequality $\left| \frac{p}{q} - x \right| < \psi(q)$ has no rational solution p/q .

Khintchine also showed that if the series $\sum_{q \geq 1} q^{-2} \psi(q)$ converges, then the set $K(\psi)$ has zero Lebesgue measure. We show that, for almost every real number x , there is a function ψ which satisfies good "regularity" conditions (for instance $q \mapsto q^2 \psi(q)$ is non-increasing),

such that the series $\sum_{q \geq 1} q^{-\psi(q)}$ converges but the inequality $\left| \frac{p}{q} - \alpha \right| < \psi(q)$ has infinitely many rational solutions $p=q$.

We also compute Hausdorff dimensions of sets of exceptions to our results (in terms of the regularity conditions on ψ):

Theorem 3.10. *Let α be a real number in the interval $[1;2]$. We denote X the set of irrational numbers such that, for every function ψ with $q^{-\psi(q)}$ non-increasing and satisfying*

$$\sum_{q=1}^{\infty} q^{-\psi(q)} = +\infty;$$

the inequality

$$\left| \frac{p}{q} - \alpha \right| < \psi(q)$$

has infinitely many rational solutions. We proved that X_1 is empty and that, for $\alpha \in (1;2]$, the Hausdorff dimension of X is $\alpha-2$ (and so, when α varies, assume all values in the interval $(1-2;1]$).

Theorem 3.11. *Let b be a positive real number. We denote Y_b as the set of the irrational numbers such that, for every function ψ with $q^{-\psi(q) \log \log q}$ non-increasing and satisfying*

$$\sum_{q=1}^{\infty} q^{-\psi(q)} = +\infty;$$

the inequality

$$\left| \frac{p}{q} - \alpha \right| < \psi(q)$$

has infinitely many rational solutions. Then the Hausdorff dimension of Y_b is $\frac{1}{1+e^{1/b}}$ (and so, when b varies, assume all values in the interval $(0;1-2)$).

One of the tools we used in the proof of this last theorem is the results from [18]: given $b; c > 1$ define $\tilde{\omega}(b; c) = \{ \alpha \in [0;1] : a_n = c^{b^n} \text{ for every } n \geq 1 \text{ and } \alpha = \sum_{n=1}^{\infty} \frac{a_n}{c^{b^n}} \}$. Then $HD(\tilde{\omega}(b; c)) = HD(\omega(b; c)) = \frac{1}{1+b}$. In one direction, a more precise result is proved in [10], according to which the Hausdorff dimension of the set

$$K := \{ \alpha \in [0;1] : \exp(b^n) \leq a_n \leq 3 \exp(b^n) \text{ for every } n \geq 1 \}$$

is equal to $\frac{1}{1+b}$. We notice that an older result from [11] states that the Hausdorff dimension of the set $\tilde{\omega}(b; c) = \{ \alpha \in [0;1] : \lim_{n \rightarrow \infty} a_n = 1 \}$ is $1-2$.

4 Back to Dynamical Systems (and Differential Geometry)

As we have seen, the sets M and L can be defined in terms of symbolic dynamics. Inspired by these characterizations, we may associate to a dynamical system together with a real function generalizations of the Markov and Lagrange spectra as follows:

Definition 4.1. Given a map $f : X \rightarrow X$ and a function $\varphi : X \rightarrow \mathbb{R}$, we define the associated dynamical Markov and Lagrange spectra as

$$M(f; \varphi) = \overline{\text{sup}}_{n \in \mathbb{N}} \varphi(f^{-n}(x)); x \in X \text{ and } \\ L(f; \varphi) = \overline{\text{limsup}}_{n \rightarrow \infty} \varphi(f^{-n}(x)); x \in X, \text{ respectively.}$$

Given a flow $(\varphi^t)_{t \in \mathbb{R}}$ in a manifold X , we define the associated dynamical Markov and Lagrange spectra as $M(f; (\varphi^t)) = \overline{\text{sup}}_{t \in \mathbb{R}} \varphi(\varphi^{-t}(x)); x \in X$ and $L(f; (\varphi^t)) = \overline{\text{limsup}}_{t \rightarrow \infty} \varphi(\varphi^{-t}(x)); x \in X$, respectively.

In a work in collaboration with A. Cerqueira and C. Matheus ([4]), we prove the following result, which generalizes a corresponding fact in the context of the classical Markov and Lagrange spectra:

Lemma 4.2. *Let $(\varphi; f)$ be a generic pair, where $\varphi : M^2 \rightarrow M^2$ is a diffeomorphism with M^2 a hyperbolic set for φ and $f : M \rightarrow \mathbb{R}$ is C^2 . Let $\varphi_s; \varphi_u$ be the projections of the horseshoe M^2 to the stable and unstable regular Cantor sets $K^s; K^u$ associated to it (along the unstable and stable foliations of M^2). Given $t \in \mathbb{R}$, we define*

$$K_t^s = \bigcap_{m \in \mathbb{Z}} \varphi^{-m}(f \circ \varphi^{-1} \circ \varphi^m)^{-1}(t); \\ K_t^u = \varphi_s^{-1}(t); K_t^u = \varphi_u^{-1}(t);$$

Then the functions $d_s(t) = HD(K_t^s)$ and $d_u(t) = HD(K_t^u)$ are continuous and coincide with the corresponding box dimensions.

The following result is a consequence of the scale recurrence lemma of [39] (see [28]):

Lemma 4.3. *Let $(\varphi; f)$ be a generic pair, where $\varphi : M^2 \rightarrow M^2$ is a diffeomorphism with M^2 a hyperbolic set for φ and $f : M \rightarrow \mathbb{R}$ is C^2 . Then*

$$HD(f(\varphi)) = \min(HD(\varphi); 1);$$

Moreover, if $HD(\varphi) > 1$ then $f(\varphi)$ has persistently non-empty interior.

Using the previous lemmas we prove a generalization of the results on dimensions of the dynamical spectra:

Theorem 4.4. ([4]) *Let $(\varphi; f)$ be a generic pair, where $\varphi : M^2 \rightarrow M^2$ is a conservative diffeomorphism with M^2 a hyperbolic set for φ and $f : M \rightarrow \mathbb{R}$ is C^2 . Then*

$$HD(L(f; \varphi) \setminus (\varphi^{-1}; t)) = HD(M(f; \varphi) \setminus (\varphi^{-1}; t)) =: d(t)$$

is a continuous real function whose image is $[0; \min(HD(\varphi); 1)]$.

(here we use the notation $L(f; \varphi) := L(f; \varphi_{\Delta})$).

In [14], P. Hubert, L. Marchese and C. Ulcigrai introduced in a similar way to the above generalizations the Lagrange spectra of closed-invariant loci for the action of $SL(2; \mathbb{R})$ on the moduli space of translation surfaces, in the context of Teichmüller dynamics, and proved that several of these spectra contain a Hall's ray.

Moreira and Romana prove the following result on Markov and Lagrange spectra for horseshoes:

Theorem 4.5 ([34]). Let $(f; \gamma)$ be a horseshoe associated to a C^2 -diffeomorphism f such that $HD(f; \gamma) > 1$. Then there is, arbitrarily close to $(f; \gamma)$ a diffeomorphism f_0 and a C^2 -neighborhood W of f_0 such that, if $(f; \gamma)$ denotes the continuation of $(f_0; \gamma_0)$ associated to $(f; \gamma) \in W$, there is an open and dense set $H \subset C^1(M; \mathbb{R})$ such that for all $f \in H$, we have

$$\text{int } L(f; \gamma) \neq \emptyset; \text{ and } \text{int } M(f; \gamma) \neq \emptyset;$$

where $\text{int } A$ denotes the interior of A .

Recently, D. Lima proved that, for typical pairs $(f; \gamma)$ as in the above theorem,

$$\sup_{f \in H} \inf_{t \in \mathbb{R}} HD(M(f; \gamma) \setminus (1; t)) < 1g = \inf_{f \in H} \inf_{t \in \mathbb{R}} \text{int}(L(f; \gamma) \setminus (1; t)) \neq \emptyset; g;$$

and Moreira ([29]) proved that, for typical pairs $(f; \gamma)$ as above, the minima of the corresponding Lagrange and Markov dynamical spectra coincide and are given by the image of a periodic point of the dynamics by the real function, solving a question by Yoccoz.

The classical Markov and Lagrange spectra can also be characterized as sets of maximum heights and asymptotic maximum heights, respectively, of geodesics in the modular surface $N = \mathbb{H}^2 / PSL(2; \mathbb{Z})$. Moreira and Romana extend in [35] the fact that these spectra have non-empty interior to the context of negative, non necessarily constant curvature as follows:

Theorem 4.6. ([35]) Let M provided with a metric g_0 be a complete noncompact surface M with finite Gaussian volume and Gaussian curvature bounded between two negative constants, i.e., if K_M denotes the Gaussian curvature, then there are constants $a; b > 0$ such that

$$a^2 \leq K_M \leq b^2 < 0;$$

Denote by SM its unitary tangent bundle and by γ its geodesic flow.

Then there is a metric g close to g_0 and a dense and C^2 -open subset $H \subset C^2(SM; \mathbb{R})$ such that

$$\text{int } M(f; g) \neq \emptyset; \text{ and } \text{int } L(f; g) \neq \emptyset;$$

for any $f \in H$, where g is the vector field defining the geodesic flow of the metric g .

Moreover, if X is a vector field sufficiently close to g then

$$\text{int } M(f; X) \neq \emptyset; \text{ and } \text{int } L(f; X) \neq \emptyset;$$

for any $f \in H$.

Recently, in collaboration with Pacifico and Romana ([36]), we proved an analogous result for Lorenz flows. Romana also proved ([59]) a corresponding result for Anosov flows in compact 3-dimensional manifolds.

Combining the techniques of this result with those of the above results in collaboration with A. Cerqueira and C. Matheus, Cerqueira, Moreira and Romana proved the following

Theorem 4.7. ([5]) Let $(f; \gamma)$ be a generic pair, where $f : M^2 \rightarrow M^2$ is a conservative diffeomorphism with $\Lambda \subset M^2$ a hyperbolic set for f and $f : M \rightarrow \mathbb{R}$ is C^2 . Then

$$HD(L(f; \gamma) \setminus (1; t)) = HD(M(f; \gamma) \setminus (1; t)) =: d(t)$$

is a continuous real function whose image is $[0; \min(HD(f; \gamma); 1)]$.

References

- [1] Y. Bugeaud, C. G. Moreira Sets of exact approximation order by rational numbers III. *Acta Arithmetica* 146 (2011), no. 2, pp. 177{193.
- [2] Y. Bugeaud, C. G. Moreira Variations autour d'un thorme mtrique de Khintchine. (French) [Variations on a metric theorem of Khintchine] *Bull. Soc. Math. France* 144 (2016), no. 3, pp. 507-538.
- [3] T. Cusick and M. Flahive, *The Marko and Lagrange spectra*, Mathematical Surveys and Monographs, 30. American Mathematical Society, Providence, RI, 1989. x+97 pp.
- [4] A. Cerqueira, C. Matheus and C. G. Moreira Continuity of Hausdor dimension across generic dynamical Lagrange and Markov spectra. <https://arxiv.org/abs/1602.04649>
- [5] A. Cerqueira, C. G. Moreira, S. Романа Continuity of Hausdor Dimension Across Generic Dynamical Lagrange and Markov Spectra II. <https://arxiv.org/abs/1711.03851>
- [6] S. Crovisier and E. Pujals Essential hyperbolicity and homoclinic bifurcations: a dichotomy phenomenon/mechanism for diffeomorphisms. *Invent. Math.* 201 (2015), no. 2, pp. 385-517.
- [7] G. A. Freiman, *Noncoincidence of the Marko and Lagrange spectra*, *Mat. Zametki* 3 (1968), 195{200; English transl., *Math. Notes* 3 (1968),125{128.
- [8] G. A. Freiman, *The initial point of Hall's ray*, *Number-theoretic studies in the Markov spectrum and in the structural theory of set addition*, pp. 87{120, 121{125. Kalinin. Gos. Univ., Moscow, 1973.
- [9] G. A. Freiman, *Diophantine approximation and geometry of numbers (The Marko spectrum)*, Kalinin. Gos. Univ., Moscow, 1975.
- [10] D. J. Feng, J. Wu, J.-C. Liang and S. Tseng, Appendix to the paper by T. Luczak - a simple proof of the lower bound: "On the fractional dimension of sets of continued fractions", *Mathematika* 44 (1997), pp. 54{55.
- [11] I. G. Good, The fractional dimensional theory of continued fractions, *Proc. Camb. Phil. Soc.* 37 (1941), pp. 199{228.
- [12] M. Hall, On the sum and product of continued fractions, *Annals of Math.*, Vol. 48 (1947), pp. 966{993.
- [13] S. Hayashi. Connecting invariant manifolds and the solution of the C^1 stability and -stability conjectures for flows. *Annals of Math.*, Vol. 145 (1997), pp. 81-137.
- [14] P. Hubert, L. Marchese, C. Ulcigrai, Lagrange spectra in Teichmller dynamics via renormalization. *Geom. Funct. Anal.* 25 (2015), no. 1, pp. 180-255.

- [15] M. Hochman and P. Shmerkin, Local entropy averages and projections of fractal measures. *Annals of Math.* 175(3):1001-1059 (2012).
- [16] O. Jenkinson and M. Pollicott, *Computing the dimension of dynamically defined sets: E_2 and bounded continued fractions*, *Ergodic Theory Dynam. Systems* 21 (2001), no. 5, 1429-1445.
- [17] O. Jenkinson and M. Pollicott, *Rigorous effective bounds on the Hausdorff dimension of continued fraction Cantor sets: a hundred decimal digits for the dimension of E_2* , Preprint (2016) available at arXiv:1611.09276.
- [18] T. Luczak On the fractional dimension of sets of continued fractions, *Mathematika* 44 (1997), pp. 50-53.
- [19] Y. Lima and C.G. Moreira, A combinatorial proof of Marstrand's theorem for products of regular Cantor sets. *Expo. Math.* 29 (2011), no. 2, 231-239.
- [20] Y. Lima and C.G. Moreira, Yet another proof of Marstrand's theorem. *Bulletin of the Brazilian Mathematical Society* 42 (2011), no. 2, 331-345.
- [21] R. Mañé, A proof of the C^1 stability conjecture. *Publ. Math. I.H.E.S.* 66 (1988), pp. 161-210.
- [22] C. Matheus, The Lagrange and Markov spectra from the dynamical point of view. <https://arxiv.org/abs/1703.01748>
- [23] J.M. Marstrand, Some fundamental geometrical properties of plane sets of fractional dimensions, *Proceedings of the London Mathematical Society* 3 (1954), vol. 4, 257-302.
- [24] A. Markov, Sur les formes quadratiques binaires infinies, *Math. Ann.* v. 15, p. 381-406, 1879.
- [25] C.G. Moreira, Stable intersections of Cantor sets and homoclinic bifurcations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 13 (1996), no. 6, pp. 741-781.
- [26] C.G. Moreira, There are no C^1 -stable intersections of regular Cantor sets. *Acta Mathematica* 206 (2011), no. 2, 311-323 .
- [27] C.G. Moreira, Geometric properties of the Markov and Lagrange spectra. <https://arxiv.org/abs/1612.05782>
- [28] C.G. Moreira, Geometric properties of images of cartesian products of regular Cantor sets by differentiable real maps. <https://arxiv.org/abs/1611.00933>
- [29] C.G. Moreira, On the minima of Markov and Lagrange Dynamical Spectra. <https://arxiv.org/abs/1711.01565>
- [30] C. Matheus, C.G. Moreira, $HD(M \cap L) > 0.353$. <https://arxiv.org/abs/1703.04302>

- [31] C. Matheus, C.G. Moreira, $HD(M \times_n L) < 0.986927$.
<https://arxiv.org/abs/1708.06258>
- [32] C. Matheus, C.G. Moreira, E. Pujals, Axiom A versus Newhouse phenomena for Benedicks-Carleson toy models. *Annales Scientifiques de l'Ecole Normale Supérieure* 46 (2013), fascicule 6, p. 857-878, 2013.
- [33] C.G. Moreira, J. Palis, M. Viana Homoclinic tangencies and fractal invariants in arbitrary dimension/Tangences homoclines et invariants fractaux en dimension arbitraire. *Comptes Rendus de l'Académie des Sciences. Série 1, Mathématique*, v. 333, p. 475-480, 2001.
- [34] C.G. Moreira, S. Romana, On the Lagrange and Markov Dynamical Spectra. *Ergodic Theory and Dynamical Systems*, pp. 1{22. doi: 10.1017/etds.2015.121.
- [35] C.G. Moreira, S. Romana, On the Lagrange and Markov Dynamical Spectra for Geodesic Flows in Surfaces with Negative Curvature. <http://arxiv.org/abs/1505.05178>
- [36] C.G. Moreira, M. J. Pacifico, S. Romana, Hausdorff Dimension, Lagrange and Markov Dynamical Spectra for Geometric Lorenz Attractors. <https://arxiv.org/abs/1611.01174>
- [37] C.G. Moreira, W. Silva, On the geometry of horseshoes in higher dimensions, <http://arxiv.org/abs/1210.2623>
- [38] L. Mora and M. Viana, Abundance of strange attractors. *Acta Math.* 171 (1993), no. 1, pp. 1-71.
- [39] C.G. Moreira and J.-C. Yoccoz Stable intersections of regular Cantor sets with large Hausdorff dimensions. *Annals Of Mathematics*, v. 154, n. 1, p. 45-96, 2001.
- [40] C.G. Moreira and J.-C. Yoccoz Tangences homoclines stables pour des ensembles hyperboliques de grande dimension fractale. *Annales Scientifiques de l'Ecole Normale Supérieure*, 43, fascicule 1, p. 1-68, 2010.
- [41] S. Newhouse, Non density of Axiom A(a) on S^2 , *Proc. A.M.S. Symp. Pure Math.*, Vol. 14, (1970), pp. 191{202.
- [42] S. Newhouse, Diffeomorphisms with infinitely many sinks, *Topology*, Vol. 13, (1974), pp. 9{18.
- [43] S. Newhouse, The abundance of wild hyperbolic sets and nonsmooth stable sets for diffeomorphisms, *Publ. Math. IHES*, Vol. 50, (1979), pp. 101{151.
- [44] J. Palis, Homoclinic orbits, hyperbolic dynamics and fractional dimension of Cantor sets, *Contemporary Mathematics* 58, (1987), pp. 203{216.
- [45] J. Palis, Homoclinic bifurcations, sensitive chaotic dynamics and strange attractors, *Dynamical Syst. and Related Topics*. World Scientific, (1991), pp. 466-473.

- [46] J. Palis and F. Takens, Cycles and measure of bifurcation sets for two-dimensional diffeomorphisms, *Invent. Math.*, Vol. 82, (1985), pp. 379{442.
- [47] J. Palis and F. Takens, Hyperbolicity and the creation of homoclinic orbits, *Annals of Math.*, Vol. 125, (1987), pp. 337{374.
- [48] J. Palis and F. Takens, Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations: fractal dimensions and infinitely many attractors, Cambridge Univ. Press, 1992.
- [49] J. Palis and J.C. Yoccoz, Rigidity of centralizers of diffeomorphisms, *Ann. Sci. cole Norm. Sup.*, Vol. 22 (1989), pp. 81{98.
- [50] J. Palis and J.C. Yoccoz, Centralizers of Anosov diffeomorphisms on tori, *Acta Mathematica*, Vol. 22 (1989), pp. 99{108.
- [51] J. Palis and J.C. Yoccoz, Differentiable conjugacies of Morse-Smale diffeomorphisms, *Bol. Soc. Brasil. Mat.*, Vol. 20 (1990), pp. 25{48.
- [52] J. Palis and J.C. Yoccoz, Homoclinic Tangencies for Hyperbolic sets of large Hausdorff Dimension Bifurcations, *Acta Mathematica*, Vol. 172 (1994), pp. 91{136.
- [53] J. Palis and J.C. Yoccoz, On the arithmetic sum of regular Cantor sets, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, Vol. 14 (1997), no.4, pp. 439{456.
- [54] J. Palis and J.C. Yoccoz, Implicit formalism for affine-like maps and parabolic composition. *Global analysis of dynamical systems*, Inst. Phys., Bristol (2001), pp. 67{87.
- [55] J. Palis and J.C. Yoccoz, Fers a cheval non uniformément hyperboliques engendrés par une bifurcation homocline et densité nulle des attracteurs / Non-uniformly hyperbolic horseshoes generated by homoclinic bifurcations and zero density of attractors, *C. R. Acad. Sci. Paris Sr. I Math.*, Vol. 333 (2001), no. 9, pp. 867-871.
- [56] J. Palis and J.C. Yoccoz, Non-uniformly hyperbolic horseshoes arising from bifurcations of Poincaré heteroclinic cycles, *Publ. Math. Inst. Hautes études Sci.*, Vol. 110 (2009), pp. 1-217.
- [57] H. Poincaré, *Les Methodes Nouvelles de la Mecanique Celeste*, III, Gauthiers-Villars, 1899.
- [58] E. Pujals and M. Sambarino, Homoclinic tangencies and hyperbolicity for surface diffeomorphisms. *Annals of Math.* (2) 151 (2000), no. 3, pp. 961-1023.
- [59] S. Romana, On the Lagrange and Markov Dynamical Spectra for Anosov Flows in dimension 3, <https://arxiv.org/abs/1605.01783>
- [60] P. Shmerkin, Moreira's Theorem on the arithmetic sum of dynamically defined Cantor sets, <http://arxiv.org/abs/0807.3709>

- [61] J.C. Yoccoz, Une erreur feconde du mathematicien Henri Poincare, *Gaz. Math.*, Vol. 107 (2006), pp. 19-26.
- [62] J.C. Yoccoz, A proof of Jakobson's theorem. https://www.college-de-france.fr/media/jean-christophe-yoccoz/UPL7416254474776698194_Jakobson_jcy.pdf